

Used for cutting tools due to being very hard.

<p><i>Each carbon atom is bonded to four others</i></p>	Very hard.	Rigid structure.
	Very high melting point.	Strong covalent bonds.
	Does not conduct electricity.	No delocalised electrons.

<p><i>Each carbon atom is bonded to three others forming layers of hexagonal rings with no covalent bonds between the layers</i></p>	Slippery.	Layers can slide over each other.
	Very high melting point.	Strong covalent bonds.
	Does conduct electricity.	Delocalised electrons between layers.

**EDEXCEL KEY CONCEPTS**  
3

**Diamond**

Graphene		Excellent conductor.	Contains delocalised electrons.
	<i>Single layer of graphite one atom thick</i>	Very strong.	Contains strong covalent bonds.

Graphene and fullerenes

<b>Graphite</b>		
Diamond, graphite, silicon dioxide	<i>Very high melting points</i>	Lots of energy needed to break strong, covalent bonds.
Used for electrodes as is inert.		

Calculations involving masses

Fullerenes		Hexagonal rings of carbon atoms with hollow shapes. Can also have rings of five (pentagonal) or seven (heptagonal) carbon atoms.
	Buckminsterfullerene, C <sub>60</sub> First fullerene to be discovered.	

The balancing numbers in a symbol equation can be calculated from the masses of reactants and products

*Convert the masses in grams to amounts in moles and convert the number of moles to simple whole number ratios.*

M <sub>r</sub>	<i>The sum of the relative atomic masses of the atoms in the numbers shown in the formula</i>	The sum of the M <sub>r</sub> of the reactants in the quantities shown equals the sum of the M <sub>r</sub> of the products in the quantities shown.	2Mg + O <sub>2</sub> → 2MgO 48g + 32g = 80g 80g = 80g

Chemical equations show the number of moles reacting and the number of moles made

**Mg + 2HCl → MgCl<sub>2</sub> + H<sub>2</sub>**

*One mole of magnesium reacts with two moles of hydrochloric acid to make one mole of magnesium chloride and one mole of hydrogen*

If you have a 60g of Mg, what mass of HCl do you need to convert it to MgCl<sub>2</sub>?

A<sub>r</sub> : Mg =24 so mass of 1 mole of Mg = 24g  
M<sub>r</sub> : HCl (1 + 35.5) so mass of 1 mole of HCl = 36.5g

So 60g of Mg is 60/24 = 2.5 moles

Balanced symbol equation tells us that for every one mole of Mg, you need two moles of HCl to react with it.

So you need 2.5x2 = 5 moles of HCl

You will need 5 x 36.5g of HCl= 182.5g

Conservation of mass	<i>No atoms are lost or made during a chemical reaction</i>	Mass of the products equals the mass of the reactants.
----------------------	---	--

Balanced symbol equations	<i>Represent chemical reactions and have the same number of atoms of each element on both sides of the equation</i>	H <sub>2</sub> + Cl <sub>2</sub> → 2HCl
	Subscript numbers show the number of atoms of the element to its left. Normal script numbers show the number of molecules.	

Avogadro constant	<i>One mole of any substance will contain the same number of particles, atoms, molecules or ions.</i>	6.02 x 10 <sup>23</sup> per mole One mole of H <sub>2</sub> O will contain 6.02 x 10 <sup>23</sup> molecules One mole of NaCl will contain 6.02 x 10 <sup>23</sup> Na <sup>+</sup> ions
-------------------	---	---

Measured in mass per given volume of solution (g/dm <sup>3</sup> )	<i>Conc. = mass (g) / volume (dm<sup>3</sup>)</i>	HT only Greater mass = higher concentration. Greater volume = lower concentration.
--	---	--