



Topic/Skill	Definition/Tips	Example						
1. Grouped Data	Data that has been <b>bundled in to categories</b> .  Seen in grouped frequency tables, histograms, cumulative frequency etc.	<table border="1"> <thead> <tr> <th>Foot length, <math>l</math>, (cm)</th> <th>Number of children</th> </tr> </thead> <tbody> <tr> <td><math>10 \leq l &lt; 12</math></td> <td>5</td> </tr> <tr> <td><math>12 \leq l &lt; 17</math></td> <td>53</td> </tr> </tbody> </table>	Foot length, $l$ , (cm)	Number of children	$10 \leq l < 12$	5	$12 \leq l < 17$	53
Foot length, $l$ , (cm)	Number of children							
$10 \leq l < 12$	5							
$12 \leq l < 17$	53							
2. Mean	<b>Add</b> up the values and <b>divide</b> by how many values there are.	The mean of 3, 4, 7, 6, 0, 4, 6 is $\frac{3 + 4 + 7 + 6 + 0 + 4 + 6}{7} = 5$						
3. Median Value	The <b>middle</b> value.  Put the data in order and find the middle one. If there are <b>two middle values</b> , find the number half way between them by <b>adding them together and dividing by 2</b> .	Find the median of: 4, 5, 2, 3, 6, 7, 6  Ordered: 2, 3, 4, <b>5</b> , 6, 6, 7  Median = 5						
4. Mode /Modal Value	<b>Most</b> frequent/common.  Can have more than one mode (called bi-modal or multi-modal) or no mode (if all values appear once)	Find the mode: 4, 5, 2, 3, 6, 4, 7, 8, 4  Mode = 4						
5. Range	<b>Highest value subtract the Smallest value</b>  Range is a 'measure of spread'. The smaller the range the more <u>consistent</u> the data.	Find the range: 3, 31, 26, 102, 37, 97.  Range = 102-3 = 99						
6. Outlier	A value that ' <b>lies outside</b> ' most of the other values in a set of data. An outlier is <b>much smaller or much larger</b> than the other values in a set of data.	<p>The scatter plot shows a positive linear trend with a line of best fit. The x-axis ranges from 0 to 100, and the y-axis ranges from 0 to 12. Most data points follow the line, but one point at approximately (25, 9) is significantly above the line and is labeled 'Outlier' with a red arrow.</p>						



Topic/Skill	Definition/Tips	Example																																						
1. Frequency Table	A record of <b>how often each value</b> in a set of data <b>occurs</b> .	<table border="1"> <thead> <tr> <th>Number of marks</th> <th>Tally marks</th> <th>Frequency</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>       </td> <td>7</td> </tr> <tr> <td>2</td> <td>    </td> <td>5</td> </tr> <tr> <td>3</td> <td>      </td> <td>6</td> </tr> <tr> <td>4</td> <td>    </td> <td>5</td> </tr> <tr> <td>5</td> <td>   </td> <td>3</td> </tr> <tr> <td><b>Total</b></td> <td></td> <td><b>26</b></td> </tr> </tbody> </table>	Number of marks	Tally marks	Frequency	1		7	2		5	3		6	4		5	5		3	<b>Total</b>		<b>26</b>																	
Number of marks	Tally marks	Frequency																																						
1		7																																						
2		5																																						
3		6																																						
4		5																																						
5		3																																						
<b>Total</b>		<b>26</b>																																						
2. Bar Chart	Represents data as vertical blocks.  $x$ – <b>axis</b> shows the <b>type</b> of data $y$ – <b>axis</b> shows the <b>frequency</b> for each type of data Each bar should be the <b>same width</b> There should be <b>gaps</b> between each bar Remember to <b>label</b> each axis.	<table border="1"> <caption>Data for Bar Chart: Frequency of Pets Owned</caption> <thead> <tr> <th>Number of pets owned</th> <th>Frequency</th> </tr> </thead> <tbody> <tr><td>0</td><td>3</td></tr> <tr><td>1</td><td>8</td></tr> <tr><td>2</td><td>12</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>2</td></tr> </tbody> </table>	Number of pets owned	Frequency	0	3	1	8	2	12	3	1	4	2																										
Number of pets owned	Frequency																																							
0	3																																							
1	8																																							
2	12																																							
3	1																																							
4	2																																							
3. Types of Bar Chart	<p><b>Compound/Composite</b> Bar Charts show data stacked on top of each other.</p> <p><b>Comparative/Dual</b> Bar Charts show data side by side.</p>	<table border="1"> <caption>Data for Compound Bar Chart: Weight (gm)</caption> <thead> <tr> <th>Sample</th> <th>Aluminum</th> <th>Carbon</th> <th>Iron</th> <th>Total</th> </tr> </thead> <tbody> <tr><td>A</td><td>25</td><td>20</td><td>15</td><td>60</td></tr> <tr><td>B</td><td>20</td><td>15</td><td>10</td><td>45</td></tr> <tr><td>C</td><td>25</td><td>20</td><td>25</td><td>70</td></tr> </tbody> </table> <table border="1"> <caption>Data for Dual Bar Chart: Rainfall (cm)</caption> <thead> <tr> <th>Month</th> <th>London</th> <th>Bristol</th> </tr> </thead> <tbody> <tr><td>Jan</td><td>15</td><td>12</td></tr> <tr><td>Feb</td><td>20</td><td>18</td></tr> <tr><td>Mar</td><td>32</td><td>35</td></tr> <tr><td>Apr</td><td>40</td><td>45</td></tr> <tr><td>May</td><td>48</td><td>50</td></tr> </tbody> </table>	Sample	Aluminum	Carbon	Iron	Total	A	25	20	15	60	B	20	15	10	45	C	25	20	25	70	Month	London	Bristol	Jan	15	12	Feb	20	18	Mar	32	35	Apr	40	45	May	48	50
Sample	Aluminum	Carbon	Iron	Total																																				
A	25	20	15	60																																				
B	20	15	10	45																																				
C	25	20	25	70																																				
Month	London	Bristol																																						
Jan	15	12																																						
Feb	20	18																																						
Mar	32	35																																						
Apr	40	45																																						
May	48	50																																						
4. Pictogram	Uses <b>pictures</b> or symbols to <b>show the value</b> of the data.  A pictogram must have a <b>key</b> .	<p>Black    🚗 🚗 🚗</p> <p>Red      🚗 🚗 🚗</p> <p>Green    🚗</p> <p>Others   🚗 🚗 🚗 🚗</p> <p>🚗 = 4 cars</p>																																						
5. Line Graph	A graph that uses <b>points connected by straight lines</b> to show how data changes in values.  This can be used for <b>time series data</b> , which is a series of data points spaced over uniform time intervals in <b>time order</b> .	<table border="1"> <caption>Data for Line Graph</caption> <thead> <tr> <th>Interval</th> <th>Value</th> </tr> </thead> <tbody> <tr><td>1</td><td>5</td></tr> <tr><td>2</td><td>4</td></tr> <tr><td>3</td><td>8</td></tr> <tr><td>4</td><td>7</td></tr> <tr><td>5</td><td>12</td></tr> <tr><td>6</td><td>8</td></tr> <tr><td>7</td><td>3</td></tr> <tr><td>8</td><td>2</td></tr> <tr><td>9</td><td>4</td></tr> </tbody> </table>	Interval	Value	1	5	2	4	3	8	4	7	5	12	6	8	7	3	8	2	9	4																		
Interval	Value																																							
1	5																																							
2	4																																							
3	8																																							
4	7																																							
5	12																																							
6	8																																							
7	3																																							
8	2																																							
9	4																																							





Topic/Skill	Definition/Tips	Example
1. Integer	A <b>whole number</b> that can be positive, negative or zero.	-3, 0, 92
2. Negative Number	A number that is <b>less than zero</b> . Can be decimals.	-8, -2.5
3. Addition	To find the <b>total</b> , or <b>sum</b> , of two or more numbers.  'add', 'plus', 'sum'	$3 + 2 + 7 = 12$
4. Subtraction	To find the <b>difference</b> between two numbers. To find out how many are left when some are taken away.  'minus', 'take away', 'subtract'	$10 - 3 = 7$
5. Multiplication	Can be thought of as <b>repeated addition</b> .  'multiply', 'times', 'product'	$3 \times 6 = 6 + 6 + 6 = 18$
6. Division	Splitting into equal parts or groups. The process of calculating the <b>number of times one number is contained within another one</b> .  'divide', 'share'	$20 \div 4 = 5$  $\frac{20}{4} = 5$
7. Remainder	The amount ' <b>left over</b> ' after dividing one integer by another.	The remainder of $20 \div 6$ is 2, because 6 divides into 20 exactly 3 times, with 2 left over.
8. BIDMAS	An acronym for the <b>order</b> you should do calculations in.  BIDMAS stands for ' <b>Brackets, Indices, Division, Multiplication, Addition and Subtraction</b> '.  Indices are also known as 'powers' or 'orders'.  With strings of division and multiplication, or strings of addition and subtraction, and no brackets, work from left to right.	$6 + 3 \times 5 = 21$ , <i>not</i> 45  $5^2 = 25$ , where the 2 is the index/power.  $12 \div 4 \div 2 = 1.5$ , <i>not</i> 6



<b>Topic/Skill</b>	<b>Definition/Tips</b>	<b>Example</b>
1. Multiple	The result of multiplying a number by an integer. The <b>times tables</b> of a number.	The first five multiples of 7 are:  7, 14, 21, 28, 35
2. Factor	A number that <b>divides exactly</b> into another number without a remainder.  It is useful to write factors in pairs	The factors of 18 are: 1, 2, 3, 6, 9, 18  The factor pairs of 18 are: 1, 18 2, 9 3, 6
3. Lowest Common Multiple (LCM)	The <b>smallest</b> number that is in the <b>times tables</b> of each of the numbers given.	The LCM of 3, 4 and 5 is 60 because it is the smallest number in the 3, 4 and 5 times tables.
4. Highest Common Factor (HCF)	The <b>biggest</b> number that <b>divides exactly</b> into two or more numbers.	The HCF of 6 and 9 is 3 because it is the biggest number that divides into 6 and 9 exactly.
5. Prime Number	A number with <b>exactly two factors</b> .  A number that can only be divided by itself and one.  The number <b>1 is not prime</b> , as it only has one factor, not two.	The first ten prime numbers are:  2, 3, 5, 7, 11, 13, 17, 19, 23, 29



Topic/Skill	Definition/Tips	Example
1. Square Number	The number you get when you <b>multiply a number by itself</b> .	<b>1, 4, 9, 16, 25, 36, 49, 64, 81, 100, 121, 144, 169, 196, 225...</b> $9^2 = 9 \times 9 = 81$
2. Square Root	The <b>number you multiply by itself</b> to get another number.  The reverse process of squaring a number.	$\sqrt{36} = 6$  because $6 \times 6 = 36$



Topic/Skill	Definition/Tips	Example
1. Expression	A mathematical statement written using <b>symbols, numbers</b> or <b>letters</b> ,	$3x + 2$ or $5y^2$
2. Equation	A statement showing that <b>two expressions are equal</b>	$2y - 17 = 15$
3. Identity	An equation that is <b>true for all values</b> of the variables  An identity uses the symbol: $\equiv$	$2x \equiv x+x$
4. Formula	Shows the <b>relationship</b> between <b>two or more variables</b>	Area of a rectangle = length x width or $A = L \times W$
5. Simplifying Expressions	<b>Collect 'like terms'</b> .  Be careful with negatives. $x^2$ and $x$ are not like terms.	$2x + 3y + 4x - 5y + 3$ $= 6x - 2y + 3$ $3x + 4 - x^2 + 2x - 1 = 5x - x^2 + 3$
6. $x$ times $x$	The answer is $x^2$ not $2x$ .	Squaring is multiplying by itself, not by 2.
7. $p \times p \times p$	The answer is $p^3$ not $3p$	If $p=2$ , then $p^3=2 \times 2 \times 2=8$ , not $2 \times 3=6$
8. $p + p + p$	The answer is $3p$ not $p^3$	If $p=2$ , then $2+2+2=6$ , not $2^3 = 8$
9. Expand	To expand a bracket, <b>multiply</b> each term <b>in the bracket</b> by the expression <b>outside</b> the bracket.	$3(m + 7) = 3m + 21$





Topic/Skill	Definition/Tips	Example
1. Inverse	<b>Opposite</b>	The inverse of addition is subtraction. The inverse of multiplication is division.
2. Writing Formulae	<b>Substitute letters for words</b> in the question.	Bob charges £3 per window and a £5 call out charge.  $C = 3N + 5$  Where N=number of windows and C=cost
3. Substitution	<b>Replace letters with numbers.</b>  Be careful of $5x^2$ . You need to square first, then multiply by 5.	$a = 3, b = 2$ and $c = 5$ . Find: 1. $2a = 2 \times 3 = 6$ 2. $3a - 2b = 3 \times 3 - 2 \times 2 = 5$ 3. $7b^2 - 5 = 7 \times 2^2 - 5 = 23$

## Topic: Accuracy



Topic/Skill	Definition/Tips	Example
1. Place Value	The <b>value</b> of where a <b>digit</b> is within a number.	In 726, the value of the 2 is 20, as it is in the 'tens' column.
2. Place Value Columns	The names of the columns that <b>determine the value of each digit</b> .  The 'ones' column is also known as the 'units' column.	<p>PLACE VALUE CHART</p> <p>Millions Hundred Thousands Ten Thousands Thousands Hundreds Tens Ones Decimal Point Tenths Hundredths Thousandths Ten-Thousandths Hundred-Thousandths Millionths</p>
3. Rounding	To make a number simpler but keep its value close to what it was.  If the <b>digit to the right</b> of the rounding digit is <b>less than 5, round down</b> . If the <b>digit to the right</b> of the rounding digit is <b>5 or more, round up</b> .	74 rounded to the nearest ten is 70, because 74 is closer to 70 than 80.  152,879 rounded to the nearest thousand is 153,000.
4. Decimal Place	The <b>position</b> of a digit to the <b>right of a decimal point</b> .	In the number 0.372, the 7 is in the second decimal place.  0.372 rounded to two decimal places is 0.37, because the 2 tells us to round down.  Careful with money - don't write £27.4, instead write £27.40
5. Estimate	To find something <b>close to the correct answer</b> .	An estimate for the height of a man is 1.8 metres.
6. Approximation	When using approximations to estimate the solution to a calculation, <b>round each number in the calculation to 1 significant figure</b> .  $\approx$ means 'approximately equal to'	$\frac{348 + 692}{0.526} \approx \frac{300 + 700}{0.5} = 2000$ <p>'Note that dividing by 0.5 is the same as multiplying by 2'</p>



Topic/Skill	Definition/Tips	Example
1. Perimeter	The <b>total distance</b> around the <b>outside</b> of a shape.  Units include: <i>mm, cm, m</i> etc.	<p style="text-align: center;">8 cm</p>  <p style="text-align: center;">5 cm</p> $P = 8 + 5 + 8 + 5 = 26cm$
2. Area	The amount of <b>space inside</b> a shape.  Units include: <i>mm<sup>2</sup>, cm<sup>2</sup>, m<sup>2</sup></i>	
3. Area of a Rectangle	<b>Length x Width</b>	 <p style="text-align: center;">9 cm</p> <p style="text-align: left;">4 cm</p> $A = 36cm^2$
4. Compound Shape	A shape made up of a <b>combination of other known shapes</b> put together.	



Topic/Skill	Definition/Tips	Example
1. Metric System	A system of measures based on: <ul style="list-style-type: none"> <li>- the metre for length</li> <li>- the kilogram for mass</li> <li>- the second for time</li> </ul> <p><b>Length: mm, cm, m, km</b>  <b>Mass: mg, g, kg</b>  <b>Volume: ml, cl, l</b></p>	$1 \text{ kilometre} = 1000 \text{ metres}$ $1 \text{ metre} = 100 \text{ centimetres}$ $1 \text{ centimetre} = 10 \text{ millimetres}$  $1 \text{ kilogram} = 1000 \text{ grams}$
2. Imperial System	A system of weights and measures originally developed in England, usually based on human quantities <p><b>Length: inch, foot, yard, miles</b>  <b>Mass: lb, ounce, stone</b>  <b>Volume: pint, gallon</b></p>	$1 \text{ lb} = 16 \text{ ounces}$ $1 \text{ foot} = 12 \text{ inches}$ $1 \text{ gallon} = 8 \text{ pints}$
3. Metric and Imperial Units	Use the <b>unitary method</b> to convert between metric and imperial units.	$5 \text{ miles} \approx 8 \text{ kilometres}$ $1 \text{ gallon} \approx 4.5 \text{ litres}$ $2.2 \text{ pounds} \approx 1 \text{ kilogram}$ $1 \text{ inch} = 2.5 \text{ centimetres}$



Topic/Skill	Definition/Tips	Example
1. Coordinates	Written in <b>pairs</b> . The <b>first</b> term is the <b>x-coordinate</b> (movement <b>across</b> ). The <b>second</b> term is the <b>y-coordinate</b> (movement <b>up or down</b> )	<p>A: (4,7) B: (-6,-3)</p>



Topic/Skill	Definition/Tips	Example
1. Fraction	A mathematical expression representing the <b>division</b> of one integer by another.  Fractions are written as <b>two numbers separated by a horizontal line.</b>	$\frac{2}{7}$ is a 'proper' fraction.  $\frac{9}{4}$ is an 'improper' or 'top-heavy' fraction.
2. Numerator	The <b>top</b> number of a fraction.	In the fraction $\frac{3}{5}$ , 3 is the numerator.
3. Denominator	The <b>bottom</b> number of a fraction.	In the fraction $\frac{3}{5}$ , 5 is the denominator.
4. Unit Fraction	A fraction where the <b>numerator is one</b> and the denominator is a positive integer.	$\frac{1}{2}, \frac{1}{3}, \frac{1}{4}$ etc. are examples of unit fractions.
5. Reciprocal	The reciprocal of a number is <b>1 divided by the number.</b>  The reciprocal of $x$ is $\frac{1}{x}$  <b>When we multiply a number by its reciprocal we get 1.</b> This is called the 'multiplicative inverse'.	The reciprocal of 5 is $\frac{1}{5}$  The reciprocal of $\frac{2}{3}$ is $\frac{3}{2}$ , because  $\frac{2}{3} \times \frac{3}{2} = 1$
6. Mixed Number	A number formed of both an <b>integer part</b> and a <b>fraction part.</b>	$3\frac{2}{5}$ is an example of a mixed number.
7. Simplifying Fractions	<b>Divide the numerator and denominator by the highest common factor.</b>	$\frac{20}{45} = \frac{4}{9}$
8. Equivalent Fractions	Fractions which represent the <b>same value.</b>	$\frac{2}{5} = \frac{4}{10} = \frac{20}{50} = \frac{60}{150} \text{ etc.}$
9. Comparing Fractions	To compare fractions, they each need to be rewritten so that they have a <b>common denominator.</b>  <b>Ascending</b> means <b>smallest to biggest.</b>  <b>Descending</b> means <b>biggest to smallest.</b>	Put in to ascending order : $\frac{3}{4}, \frac{2}{3}, \frac{5}{6}, \frac{1}{2}$ .  Equivalent: $\frac{9}{12}, \frac{8}{12}, \frac{10}{12}, \frac{6}{12}$  Correct order: $\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{5}{6}$
10. Fraction of an Amount	<b>Divide</b> by the <b>bottom</b> , <b>times</b> by the <b>top</b>	Find $\frac{2}{5}$ of £60  $60 \div 5 = 12$ $12 \times 2 = 24$
11. Adding or Subtracting Fractions	Find the <b>LCM of the denominators</b> to find a common denominator. Use equivalent fractions to change each fraction to the <b>common denominator.</b> Then just <b>add or subtract the numerators</b>	$\frac{2}{3} + \frac{4}{5}$ Multiples of 3: 3, 6, 9, 12, <b>15..</b> Multiples of 5: 5, 10, <b>15..</b> LCM of 3 and 5 = 15



	and keep the <b>denominator the same.</b>	$\frac{2}{3} = \frac{10}{15}$ $\frac{4}{4} = \frac{12}{12}$ $\frac{5}{5} = \frac{15}{15}$ $\frac{10}{15} + \frac{12}{15} = \frac{22}{15} = 1\frac{7}{15}$
12. Multiplying Fractions	<b>Multiply the numerators</b> together and <b>multiply the denominators</b> together.	$\frac{3}{8} \times \frac{2}{9} = \frac{6}{72} = \frac{1}{12}$
13. Dividing Fractions	<b>‘Keep it, Flip it, Change it – KFC’</b> Keep the first fraction the same Flip the second fraction upside down Change the divide to a multiply  Multiply by the reciprocal of the second fraction.	$\frac{3}{4} \div \frac{5}{6} = \frac{3}{4} \times \frac{6}{5} = \frac{18}{20} = \frac{9}{10}$





Topic/Skill	Definition/Tips	Example
1. Percentage	<b>Number of parts per 100.</b>	31% means $\frac{31}{100}$
2. Finding 10%	To find <b>10%</b> , <b>divide by 10</b>	10% of £36 = $36 \div 10 = £3.60$
3. Finding 1%	To find <b>1%</b> , <b>divide by 100</b>	1% of £8 = $8 \div 100 = £0.08$
4. Percentage Change	$\frac{\text{Difference}}{\text{Original}} \times 100\%$	A games console is bought for £200 and sold for £250.  % change = $\frac{50}{200} \times 100 = 25\%$
5. Fractions to Decimals	<b>Divide the numerator by the denominator</b> using the bus stop method.	$\frac{3}{8} = 3 \div 8 = 0.375$
6. Decimals to Fractions	<b>Write as a fraction</b> over 10, 100 or 1000 and simplify.	$0.36 = \frac{36}{100} = \frac{9}{25}$
7. Percentages to Decimals	<b>Divide by 100</b>	$8\% = 8 \div 100 = 0.08$
8. Decimals to Percentages	<b>Multiply by 100</b>	$0.4 = 0.4 \times 100\% = 40\%$
9. Fractions to Percentages	Percentage is just a fraction out of 100. <b>Make the denominator 100 using equivalent fractions.</b> When the denominator doesn't go in to 100, use a calculator and <b>multiply the fraction by 100.</b>	$\frac{3}{25} = \frac{12}{100} = 12\%$  $\frac{9}{17} \times 100 = 52.9\%$
10. Percentages to Fractions	Percentage is just a fraction out of 100. <b>Write the percentage over 100</b> and simplify.	$14\% = \frac{14}{100} = \frac{7}{50}$



Topic/Skill	Definition/Tips	Example
1. Probability	<p>The <b>likelihood/chance</b> of something happening.</p> <p>Is expressed as a number <b>between 0 (impossible) and 1 (certain)</b>.</p> <p>Can be expressed as a fraction, decimal, percentage or in words (likely, unlikely, even chance etc.)</p>	
2. Theoretical Probability	$\frac{\text{Number of Favourable Outcomes}}{\text{Total Number of Possible Outcomes}}$	<p>Probability of rolling a 4 on a fair 6-sided die = <math>\frac{1}{6}</math>.</p>
3. Relative Frequency	$\frac{\text{Number of Successful Trials}}{\text{Total Number of Trials}}$	<p>A coin is flipped 50 times and lands on Tails 29 times.</p> <p>The relative frequency of getting Tails = <math>\frac{29}{50}</math>.</p>
4. Expected Outcomes	<p>To find the number of expected outcomes, <b>multiply</b> the <b>probability</b> by the <b>number of trials</b>.</p>	<p>The probability that a football team wins is 0.2 How many games would you expect them to win out of 40?</p> $0.2 \times 40 = 8 \text{ games}$
5. Exhaustive	<p>Outcomes are <b>exhaustive</b> if they <b>cover the entire range of possible outcomes</b>.</p> <p>The <b>probabilities</b> of an <b>exhaustive</b> set of outcomes <b>adds up to 1</b>.</p>	<p>When rolling a six-sided die, the outcomes 1, 2, 3, 4, 5 and 6 are exhaustive, because they cover all the possible outcomes.</p>
6. Mutually Exclusive	<p>Events are mutually exclusive if they <b>cannot happen at the same time</b>.</p> <p>The <b>probabilities</b> of an exhaustive set of <b>mutually exclusive</b> events <b>adds up to 1</b>.</p>	<p>Examples of mutually exclusive events:</p> <ul style="list-style-type: none"> <li>- Turning left and right</li> <li>- Heads and Tails on a coin</li> </ul> <p>Examples of non mutually exclusive events:</p> <ul style="list-style-type: none"> <li>- King and Hearts from a deck of cards, because you can pick the King of Hearts</li> </ul>





Topic/Skill	Definition/Tips	Example
1. Ratio	Ratio compares the size of <b>one part</b> to <b>another part</b> .  Written using the ':' symbol.	$3 : 1$ 
2. Proportion	Proportion compares the size of <b>one part</b> to the size of the <b>whole</b> .  Usually written as a fraction.	In a class with 13 boys and 9 girls, the proportion of boys is $\frac{13}{22}$ and the proportion of girls is $\frac{9}{22}$
3. Simplifying Ratios	<b>Divide</b> all parts of the ratio by a <b>common factor</b> .	$5 : 10 = 1 : 2$ (divide both by 5) $14 : 21 = 2 : 3$ (divide both by 7)
4. Ratios in the form $1 : n$ or $n : 1$	<b>Divide</b> both parts of the ratio by one of the numbers to make <b>one part equal 1</b> .	$5 : 7 = 1 : \frac{7}{5}$ in the form $1 : n$ $5 : 7 = \frac{5}{7} : 1$ in the form $n : 1$
5. Sharing in a Ratio	<b>1. Add</b> the total parts of the ratio. <b>2. Divide</b> the amount to be shared by this value to find the value of one part. <b>3. Multiply</b> this value by each part of the ratio.  Use only if you <b>know the total</b> .	Share £60 in the ratio $3 : 2 : 1$ .  $3 + 2 + 1 = 6$ $60 \div 6 = 10$ $3 \times 10 = 30, 2 \times 10 = 20, 1 \times 10 = 10$ $\pounds 30 : \pounds 20 : \pounds 10$
6. Proportional Reasoning	Comparing two things using <b>multiplicative reasoning</b> and applying this to a new situation.  Identify one multiplicative link and use this to find missing quantities.	
7. Unitary Method	Finding the <b>value of a single unit</b> and then finding the necessary value by <b>multiplying</b> the single unit value.	3 cakes require 450g of sugar to make. Find how much sugar is needed to make 5 cakes.  $3 \text{ cakes} = 450\text{g}$ So $1 \text{ cake} = 150\text{g}$ ( $\div$ by 3) So $5 \text{ cakes} = 750 \text{ g}$ ( $\times$ by 5)
8. Ratio already shared	Find what <b>one part</b> of the ratio is worth using the <b>unitary method</b> .	Money was shared in the ratio $3:2:5$ between Ann, Bob and Cat. Given that Bob had £16, found out the total amount of money shared.  $\pounds 16 = 2 \text{ parts}$ So $\pounds 8 = 1 \text{ part}$ $3 + 2 + 5 = 10 \text{ parts}$ , so $8 \times 10 = \pounds 80$
9. Best Buys	Find the <b>unit cost</b> by <b>dividing the price by the quantity</b> . The <b>lowest</b> number is the best value.	8 cakes for $\pounds 1.28 \rightarrow 16\text{p}$ each ( $\div$ by 8) 13 cakes for $\pounds 2.05 \rightarrow 15.8\text{p}$ each ( $\div$ by 13) Pack of 13 cakes is best value.



Topic/Skill	Definition/Tips	Example
1. Types of Angles	<p><b>Acute angles</b> are less than <math>90^\circ</math>.</p> <p><b>Right angles</b> are exactly <math>90^\circ</math>.</p> <p><b>Obtuse angles</b> are greater than <math>90^\circ</math> but less than <math>180^\circ</math>.</p> <p><b>Reflex angles</b> are greater than <math>180^\circ</math> but less than <math>360^\circ</math>.</p>	<p>Acute      Right      Obtuse      Reflex</p>
2. Angle Notation	<p>Can use <b>one lower-case</b> letters, eg. <math>\theta</math> or <math>x</math></p> <p>Can use <b>three upper-case</b> letters, eg. <math>BAC</math></p>	
3. Angles at a Point	<p><b>Angles around a point add up to <math>360^\circ</math>.</b></p>	<p><math>a + b + c + d = 360^\circ</math></p>
4. Angles on a Straight Line	<p><b>Angles around a point on a straight line add up to <math>180^\circ</math>.</b></p>	<p><math>x + y = 180^\circ</math></p>
5. Opposite Angles	<p><b>Vertically opposite angles are equal.</b></p>	<p><math>x = y</math></p>
6. Angles in a Triangle	<p><b>Angles in a triangle add up to <math>180^\circ</math>.</b></p>	
7. Types of Triangles	<p><b>Right Angle</b> Triangles have a <math>90^\circ</math> angle in.</p> <p><b>Isosceles</b> Triangles have <b>2 equal sides</b> and <b>2 equal base angles</b>.</p> <p><b>Equilateral</b> Triangles have <b>3 equal sides</b> and <b>3 equal angles (<math>60^\circ</math>)</b>.</p> <p><b>Scalene</b> Triangles have <b>different sides</b> and <b>different angles</b>.</p> <p><b>Base angles in an isosceles triangle are equal.</b></p>	<p>Right Angled      Isosceles</p> <p>Equilateral      Scalene</p>







8. Angles in a Quadrilateral	<b>Angles in a quadrilateral add up to <math>360^\circ</math>.</b>	
9. Regular	A shape is regular if all the <b>sides</b> and all the <b>angles</b> are equal.	






Topic/Skill	Definition/Tips	Example
1. Linear Sequence	A number pattern with a <b>common difference</b> .	2, 5, 8, 11... is a linear sequence
2. Term	<b>Each value</b> in a sequence is called a term.	In the sequence 2, 5, 8, 11..., 8 is the third term of the sequence.
3. Term-to-term rule	A rule which allows you to <b>find the next term</b> in a sequence if you <b>know the previous term</b> .	First term is 2. Term-to-term rule is 'add 3'  Sequence is: 2, 5, 8, 11...
4. nth term	A rule which allows you to <b>calculate the term</b> that is in the <b>nth position</b> of the sequence.  Also known as the 'position-to-term' rule.  <b>n</b> refers to the <b>position</b> of a term in a sequence.	nth term is $3n - 1$  The 100 <sup>th</sup> term is $3 \times 100 - 1 = 299$
5. Finding the nth term of a linear sequence	1. Find the <b>difference</b> . 2. <b>Multiply that by n</b> . 3. Substitute $n = 1$ to <b>find out what number you need to add or subtract to get the first number in the sequence</b> .	Find the nth term of: 3, 7, 11, 15...  1. Difference is +4 2. Start with $4n$ 3. $4 \times 1 = 4$ , so we need to subtract 1 to get 3. nth term = $4n - 1$
6. Triangular numbers	The sequence which comes from a pattern of dots that form a triangle.  $1, 3, 6, 10, 15, 21 \dots$	<p>1      3      6      10</p>



Topic/Skill	Definition/Tips	Example																
1. Coordinates	Written in <b>pairs</b> . The <b>first</b> term is the <b>x-coordinate</b> (movement <b>across</b> ). The <b>second</b> term is the <b>y-coordinate</b> (movement <b>up or down</b> )	 <div style="display: inline-block; vertical-align: top; margin-left: 20px;">                 A: (4,7)                  B: (-6,-3)             </div>																
2. Midpoint of a Line	<p><b>Method 1: add the x coordinates and divide by 2, add the y coordinates and divide by 2</b></p> <p>Method 2: Sketch the line and find the values half way between the two x and two y values.</p>	<p>Find the midpoint between (2,1) and (6,9)</p> $\frac{2+6}{2} = 4 \text{ and } \frac{1+9}{2} = 5$ <p>So, the midpoint is (4,5)</p>																
3. Linear Graph	<p><b>Straight line</b> graph.</p> <p>The general equation of a linear graph is <math>y = mx + c</math></p> <p>where <b>m</b> is the <b>gradient</b> and <b>c</b> is the <b>y-intercept</b>.</p> <p>The <b>equation</b> of a linear graph can contain an <b>x-term</b>, a <b>y-term</b> and a <b>number</b>.</p>	<p>Example:</p>  <div style="display: inline-block; vertical-align: top; margin-left: 20px;"> <p>Other examples:</p> <math>x = y</math>  <math>y = 4</math>  <math>x = -2</math>  <math>y = 2x - 7</math>  <math>y + x = 10</math>  <math>2y - 4x = 12</math> </div>																
4. Plotting Linear Graphs	<p><b>Method 1: Table of Values</b> Construct a table of values to calculate coordinates.</p> <p><b>Method 2: Gradient-Intercept Method</b> (use when the equation is in the form <math>y = mx + c</math>)</p> <ol style="list-style-type: none"> <li>Plots the y-intercept</li> <li>Using the gradient, plot a second point.</li> <li>Draw a line through the two points plotted.</li> </ol> <p><b>Method 3: Cover-Up Method</b> (use when the equation is in the form <math>ax + by = c</math>)</p> <ol style="list-style-type: none"> <li>Cover the x term and solve the resulting equation. Plot this on the x – axis.</li> <li>Cover the y term and solve the resulting equation. Plot this on the y – axis.</li> <li>Draw a line through the two points plotted.</li> </ol>	<table border="1" style="margin-bottom: 20px; width: 100%; text-align: center;"> <tr style="background-color: #FFD700;"> <th>x</th> <td>-3</td> <td>-2</td> <td>-1</td> <td>0</td> <td>1</td> <td>2</td> <td>3</td> </tr> <tr style="background-color: #FFD700;"> <th>y = x + 3</th> <td>0</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> <td>6</td> </tr> </table>  	x	-3	-2	-1	0	1	2	3	y = x + 3	0	1	2	3	4	5	6
x	-3	-2	-1	0	1	2	3											
y = x + 3	0	1	2	3	4	5	6											



Topic/Skill	Definition/Tips	Example
1. Congruent Shapes	Shapes are congruent if they are <b>identical - same shape and same size.</b>  Shapes can be rotated or reflected but still be congruent.	
2. Similar Shapes	Shapes are similar if they are the <b>same shape but different sizes.</b>  The proportion of the matching sides must be the same, meaning the ratios of corresponding sides are all equal.	
3. Scale Factor	The <b>ratio of corresponding sides</b> of two similar shapes.  To find a scale factor, <b>divide a length</b> on one shape <b>by the corresponding length</b> on a similar shape.	 <p style="text-align: center;">Scale Factor = <math>15 \div 10 = 1.5</math></p>
4. Finding missing lengths in similar shapes	1. Find the <b>scale factor.</b> 2. <b>Multiply or divide</b> the corresponding side to find a missing length.  If you are finding a missing length on the larger shape you will need to multiply by the scale factor.  If you are finding a missing length on the smaller shape you will need to divide by the scale factor.	 <p style="text-align: center;">Scale Factor = <math>3 \div 2 = 1.5</math>  <math>x = 4.5 \times 1.5 = 6.75cm</math></p>



Topic/Skill	Definition/Tips	Example
1. Translation	<p><b>Translate</b> means to <b>move a shape</b>. The shape does not change <b>size</b> or <b>orientation</b>.</p>	
2. Column Vector	<p>In a column vector, the <b>top</b> number moves <b>left (-) or right (+)</b> and the <b>bottom</b> number moves <b>up (+) or down (-)</b></p>	<p><math>\begin{pmatrix} 2 \\ 3 \end{pmatrix}</math> means '2 right, 3 up'  <math>\begin{pmatrix} -1 \\ -5 \end{pmatrix}</math> means '1 left, 5 down'</p>
3. Rotation	<p>The size does not change, but the <b>shape is turned around a point</b>.  Use tracing paper.</p>	<p>Rotate Shape A 90° anti-clockwise about (0,1)</p>
4. Reflection	<p>The size does not change, but the shape is '<b>flipped</b>' like in a <b>mirror</b>.</p> <p>Line <math>x = ?</math> is a <b>vertical line</b>.                      Line <math>y = ?</math> is a <b>horizontal line</b>.                      Line <math>y = x</math> is a <b>diagonal line</b>.</p>	<p>Reflect shape C in the line <math>y = x</math></p>
5. Enlargement	<p>The shape will get <b>bigger or smaller</b>. Multiply each side by the <b>scale factor</b>.</p>	<p>Scale Factor = 3 means '3 times larger = multiply by 3'                       Scale Factor = <math>\frac{1}{2}</math> means 'half the size = divide by 2'</p>

<p>6. Finding the Centre of Enlargement</p>	<p>Draw <b>straight lines</b> through <b>corresponding corners</b> of the two shapes. The centre of enlargement is the point <b>where all the lines cross over</b>.</p> <p>Be careful with negative enlargements as the corresponding corners will be the other way around.</p>	<p>A to B is an enlargement SF 2 about the point (2,1)</p>
<p>7. Describing Transformations</p>	<p>Give the following information when describing each transformation:</p> <p>Look at the number of marks in the question for a hint of how many pieces of information are needed.</p> <p>If you are asked to describe a 'transformation', you need to say the <b>name of the type of transformation</b> as well as the other details.</p>	<ul style="list-style-type: none"> <li>- Translation, Vector</li> <li>- Rotation, Direction, Angle, Centre</li> <li>- Reflection, Equation of mirror line</li> <li>- Enlargement, Scale factor, Centre of enlargement</li> </ul>